

**Table 1:** Guidelines for Recreational Diving with Diabetes - Summary Form<sup>1</sup>

<p><b>Selection and Surveillance</b></p> <ul style="list-style-type: none"> <li>• Age <math>\geq 18</math> years (<math>\geq 16</math> years if in special training program)</li> <li>• Delay diving after start/change in medication             <ul style="list-style-type: none"> <li>- 3 months with oral hypoglycemic agents (OHA)</li> <li>- 1 year after initiation of insulin therapy</li> </ul> </li> <li>• No episodes of hypoglycemia or hyperglycemia requiring intervention from a third party for at least one year</li> <li>• No history of hypoglycemia unawareness</li> <li>• <math>HbA_{1c} \leq 9\%</math> no more than one month prior to initial assessment and at each annual review             <ul style="list-style-type: none"> <li>- values <math>&gt;9\%</math> indicate the need for further evaluation and possible modification of therapy</li> </ul> </li> <li>• No significant secondary complications from diabetes</li> <li>• Physician/Diabetologist should carry out annual review and determine that diver has good understanding of disease and effect of exercise             <ul style="list-style-type: none"> <li>- in consultation with an expert in diving medicine, as required</li> </ul> </li> <li>• Evaluation for silent ischemia for candidates <math>&gt;40</math> years of age             <ul style="list-style-type: none"> <li>- after initial evaluation, periodic surveillance for silent ischemia can be in accordance with accepted local/national guidelines for the evaluation of diabetics</li> </ul> </li> <li>• Candidate documents intent to follow protocol for divers with diabetes and to cease diving and seek medical review for any adverse events during diving possibly related to diabetes</li> </ul>
<p><b>Scope of Diving</b></p> <ul style="list-style-type: none"> <li>• Diving should be planned to avoid             <ul style="list-style-type: none"> <li>- depths <math>&gt;100</math> fsw (30 msw)</li> <li>- durations <math>&gt;60</math> minutes</li> <li>- compulsory decompression stops</li> <li>- overhead environments (e.g., cave, wreck penetration)</li> <li>- situations that may exacerbate hypoglycemia (e.g., prolonged cold and arduous dives)</li> </ul> </li> <li>• Dive buddy/leader informed of diver's condition and steps to follow in case of problem</li> <li>• Dive buddy should not have diabetes</li> </ul>
<p><b>Glucose Management on the Day of Diving</b></p> <ul style="list-style-type: none"> <li>• General self-assessment of fitness to dive</li> <li>• Blood glucose (BG) <math>\geq 150</math> mg·dL<sup>-1</sup> (8.3 mmol·L<sup>-1</sup>), stable or rising, before entering the water             <ul style="list-style-type: none"> <li>- complete a minimum of three pre-dive BG tests to evaluate trends                 <ul style="list-style-type: none"> <li>▪ 60 minutes, 30 minutes and immediately prior to diving</li> </ul> </li> <li>- alterations in dosage of OHA or insulin on evening prior or day of diving may help</li> </ul> </li> <li>• Delay dive if BG             <ul style="list-style-type: none"> <li>- <math>&lt;150</math> mg·dL<sup>-1</sup> (8.3 mmol·L<sup>-1</sup>)</li> <li>- <math>&gt;300</math> mg·dL<sup>-1</sup> (16.7 mmol·L<sup>-1</sup>)</li> </ul> </li> <li>• Rescue medications             <ul style="list-style-type: none"> <li>- carry readily accessible oral glucose during all dives</li> <li>- have parenteral glucagon available at the surface</li> </ul> </li> <li>• If hypoglycemia noticed underwater, the diver should surface (with buddy), establish positive buoyancy, ingest glucose and leave the water</li> <li>• Check blood sugar frequently for 12-15 hours after diving</li> <li>• Ensure adequate hydration on days of diving</li> <li>• Log all dives (include BG test results and all information pertinent to diabetes management)</li> </ul>

<sup>1</sup> For full text see: Pollock NW, Ugucioni DM, Dear GdeL, eds. Diabetes and recreational diving: guidelines for the future. Proceedings of the UHMS/DAN 2005 June 19 Workshop. Durham, NC: Divers Alert Network; 2005.